

Technical Data Sheet

WesternFroxx Kit anti-rabbit HRP

for immunodetection

Order number: 6666

The special 10X wash buffer can be ordered separately: 5570ML500, 1 x 500 ml

WesternFroxx enables **immunodetection in one step with one solution only**. Blocking and binding of the primary and secondary antibody occur simultaneously with highest specificity and sensitivity. Blotting procedure, membranes and signal detection do not have to be modified with WesternFroxx. Proteins are detected by ECL or other HRP substrates. Despite the fast incubation process, WesternFroxx leads to clear and distinct bands suitable for publishing. The reason: WesternFroxx solution does block the membrane much faster and more effectively compared to standard blockers. Therefore, the primary and secondary antibodies don't bind to the membrane non-specifically and a faster incubation process is enabled. Furthermore, by using the special Washing buffer after blotting the membrane is released from components of the transfer buffer (e.g. methanol or ethanol) and perfectly prepared for the extremely fast blocking step. WesternFroxx is also available as anti-mouse HRP (order number 5555).

Kit components

The kits contain the WesternFroxx detection solution and a special 10X washing buffer

Solution 1 10X Washing buffer

Solution 2 WesternFroxx anti-rabbit HRP solution containing blocking reagent and secondary antibody (anti-rabbit HRP)

Preparation

Before you start with WesternFroxx

- x Transfer your proteins to the membrane of choice (PVDF, nitrocellulose) as usually.
- x Dilute the provided 10X washing buffer (Solution 1) with water.
- × Add your primary antibody into 20 ml of WesternFroxx Solution 2. The final concentration of the antibody should be 0.2 to $0.5 \mu g / ml$.



neoFroxx GmbH Marie-Curie-Str. 3 D-64683 Einhausen www.neofroxx.com Phone +49 (6251) 989 24-0 info@neofroxx.com



Protocol

- 1. Wash the membrane 3 x 5 min. while shaking with 20 ml of 1X Solution 1 (minigel).
- Incubate the membrane with WesternFroxx Solution 2 (containing your primary antibody of choice) for 15 min. to 1 hour.

Caution! The WesternFroxx solution can be reused up to 5 times. Keep it in the fridge.

- 3. Wash the membrane again for 3×5 min with 1X Solution 1.
- 4. Start the detection with ECL-Solution (we recommend ECL Xtrasensitivity Kit, order number 5560).

Related products

- 7777 WesternFroxx all-in-one Protein Ladder (15 - 200 kDa) for immunodetection 1052 Protein Ladder (6.5 - 212 kDa), unstained for molecular biology 1123 Protein Ladder (11 – 245 kDa), prestained for molecular biology 1288 Protein Ladder (10 – 180 kDa), prestained for molecular biology 5555 WesternFroxx Kit anti-mouse HRP for immunodetection 5560 ECL Xtrasensitivity Kit for immunodetection 1550 BSA blocking solution for immunodetection 1080 **CleanBlot Background Minimizer for immunodetection** 1095 Peptide blocking solution (BSA-free) for immunodetection 1080 Casein high-end blocking solution for immunodetection 1102 Acrylamide Xtra solution 30 % - Mix 37.5:1 for electrophoresis 1248 Acrylamide Xtra solution 40 % - Mix 37.5:1 for electrophoresis 1106 Acrylamide Xtra solution 30 % - Mix 29:1 for electrophoresis 1912 Coomassie® brilliant blue R-250 (C.I. 42660) for biochemistry 1277 SDS Xtrapure for biochemistry 8027 **TEMED** for biochemistry 1610 Ammonium persulfate for molecular biology
 - 1125 Tris Xtrapure for biochemistry
 - 1275 Glycine for biochemistry

JB30112018



neoFroxx GmbH Marie-Curie-Str. 3 D-64683 Einhausen www.neofroxx.com Phone +49 (6251) 989 24-0 info@neofroxx.com